



Growth parameter estimates of *Listeria monocytogenes* in cooked chicken: effect of preparation of inoculum

Birk, Tina; Smith Ottosen, Sussi; Hansen, Tina Beck

Published in:
10th International conference on predictive modelling in food

Publication date:
2017

Document Version
Publisher's PDF, also known as Version of record

[Link back to DTU Orbit](#)


Citation (APA):
Birk, T., Smith Ottosen, S., & Hansen, T. B. (2017). Growth parameter estimates of *Listeria monocytogenes* in cooked chicken: effect of preparation of inoculum. In *10th International conference on predictive modelling in food*

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.


- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.



10TH INTERNATIONAL CONFERENCE ON **PREDICTIVE MODELLING IN FOOD**

CÓRDOBA, SPAIN, 26-29 SEPTEMBER, 2017
GOVERNMENT BUILDING OF THE UNIVERSITY OF CÓRDOBA



CÓRDOBA, SEPTEMBER, 2017
ICPMF10
INTERNATIONAL CONFERENCE ON PREDICTIVE MODELLING IN FOOD



UNIVERSIDAD DE CÓRDOBA

www.icpmf10.com

P. 17.- GROWTH PARAMETER ESTIMATES OF *LISTERIA MONOCYTOGENES* IN COOKED CHICKEN: EFFECT OF PREPARATION OF INOCULUMS.

Tina Birk¹, Sussi Smith Ottosen², Tina Beck Hansen¹

¹ The National Food Institute, Technical University of Denmark, Kemitorvet, Lyngby, Denmark

² University College Zealand, Campus Roskilde, Maglegaardsvej 8, Denmark

Keywords: *Listeria*, challenge test, growth modelling

INTRODUCTION AND OBJETIVES

Estimates of lag times and growth rates from growth curves for *Listeria monocytogenes* on slices of sous-vide cooked chicken breast were compared for four inoculum preparation procedures. Effects were investigated at different temperatures between 6 and 24 °C.

MATERIAL AND METHODS

Slices of sous-vide cooked chicken breast (12 ± 2.7 g) were inoculated with five 10- μ l-spots of a four-strain cocktail of *L. monocytogenes* resulting in inoculation volumes of 0.44 ± 0.11 %. Each strain was inoculated into Brain Heart Infusion broth (BHI) and incubated at 30 °C for 24 h before sub-cultured in four different ways, mixing into a cocktail and used for inoculation; i) dilution in fresh BHI (direct30), ii) chilled incubation for 3 d before dilution in fresh BHI (coldBHI), iii) chilled incubation for 3 d before dilution in Maximum Recovery Diluent (MRD) (coldMRD) and iv) dilution in fresh BHI followed by chilled incubation for 3 d (direct8). Direct30 and direct8 procedures were tested against coldBHI at 8 and 19 °C, whereas the coldMRD procedure was compared to coldBHI in the temperature range from 6 to 24 °C. Growth curves were graphically compared and fitted to Baranyi & Roberts models using DMFit. Lag times (L) and max specific growth rates (μ_{max}) were used for statistical analyses using t-tests and for developing secondary models for the coldMRD and coldBHI procedures. Secondary models were cross-validated using bias (Bf) and accuracy (Af) factors.

RESULTS

Growth curves obtained at 19 °C showed signs of injured cells repairing during the first 3 h when the direct30 procedure was used compared to coldBHI. However, no clear statistical differences were found for L - ($P = 0.07$) or μ_{max} -values ($P = 0.50$). At 8 °C, growth curves obtained for direct30 and coldBHI were identical in the first 30 h then counts became more scattered with a tendency to have higher counts for the coldBHI procedure. Nevertheless, no statistical differences were found neither for L - ($P = 0.99$) nor μ_{max} -values ($P = 0.76$). The direct8 procedure resulted, in significantly ($P = 0.04$) longer L at 19 °C but similar μ_{max} ($P = 0.12$) when compared to coldBHI. At 8 °C, growth curves obtained using the direct8 procedure appeared to have no lag phase when fitted. However, these curves showed clear signs of injured cells repairing during the first 5 – 7 h which suggested that the obtained L -values of 0 could be an artefact resulting from the curve-fitting procedure. Based on Bf , μ_{max} -estimates obtained using the coldBHI procedure were on average 12 % higher than results observed for coldMRD. An Af of 1.17 was found when coldMRD observations were used for validating the coldBHI secondary μ_{max} -model. In the opposite case, a slightly lower Af of 1.14 was found. Graphical comparison showed approx. 3-fold longer L for coldMRD below 10 °C.

CONCLUSIONS

Even with very low inoculation volume, estimates of lag times and growth rates of *L. monocytogenes* were significantly affected by the procedure used for preparation of the inoculum.